

1 $f(x) \equiv x^3 + x^2 - 22x - 40.$

- a Show that $(x + 2)$ is a factor of $f(x)$. (2)
 b Express $f(x)$ as the product of three linear factors. (4)
 c Solve the equation $f(x) = 0$. (1)

2 $f(x) \equiv x^3 - 2x^2 + kx + 1.$

Given that the remainder when $f(x)$ is divided by $(x - 2)$ and the remainder when $f(x)$ is divided by $(x + 3)$ are equal,


- a find the value of the constant k , (4)
 b find the remainder when $f(x)$ is divided by $(x + 2)$. (2)

3 The polynomial $p(x)$ is defined by

$$p(x) \equiv 2x^3 - 9x^2 - 2x + 11.$$

- a Find the remainder when $p(x)$ is divided by $(x + 2)$. (2)
 b Find the quotient and remainder when $p(x)$ is divided by $(x - 4)$. (3)

4



The diagram shows the curve with the equation $y = x^3 - 5x^2 - 8x + 12$.

- a State the coordinates of the point A where the curve crosses the y -axis. (1)
 The curve crosses the x -axis at the points B , C and D . Given that C has coordinates $(1, 0)$,
 b find the coordinates of the points B and D . (6)

5 $f(x) \equiv x^3 - 3x^2 + kx + 8.$

Given that $(x - 1)$ is a factor of $f(x)$,

- a find the value of k , (2)
 b solve the equation $f(x) = 0$. (5)

6 Solve the equation

$$2x^3 + x^2 - 13x + 6 = 0. \quad (7)$$

7 The polynomial $p(x)$ is defined by

$$p(x) \equiv bx^3 + ax^2 - 10x + b,$$

where a and b are constants.

Given that when $p(x)$ is divided by $(x + 1)$ the remainder is 3,

- a find the value of a . (2)
 Given also that when $p(x)$ is divided by $(3x - 1)$ the remainder is -1 ,
 b find the value of b . (3)

- 8 $f(x) \equiv x^3 - 7x^2 + x + 10.$
- a Find the remainder when $f(x)$ is divided by $(x + 1).$ (2)
- b Hence, or otherwise, solve the equation $f(x) = 1,$ giving your answers in exact form. (6)
- 9 $f(x) \equiv 3x^3 + kx^2 - 7x + 2k.$
- When $f(x)$ is divided by $(3x - 2)$ the remainder is 6.
- Find the value of the constant $k.$ (3)
- 10 $f(x) \equiv 2x^3 - 7x^2 + 4x - 3.$
- a Show that $(x - 3)$ is a factor of $f(x).$ (2)
- b Hence, express $f(x)$ as the product of a linear factor and a quadratic factor. (3)
- c Show that there is only one real solution to the equation $f(x) = 0.$ (3)
- 11 The polynomial $f(x)$ is defined by
- $$f(x) \equiv x^3 + px + q,$$
- where p and q are constants.
- Given that $(x - 2)$ is a factor of $f(x),$
- a find an expression for q in terms of $p.$ (2)
- Given also that when $f(x)$ is divided by $(x + 1)$ the remainder is $-15,$
- b find the values of p and $q.$ (4)
- 12 $f(x) \equiv x^3 + 4x^2 - 9.$
- Given that $x = -3$ is a solution to the equation $f(x) = 0,$ find the other two solutions correct to 2 decimal places. (6)
- 13 $f(x) \equiv (x + k)^3 - 8.$
- Given that when $f(x)$ is divided by $(x + 2)$ the remainder is $-7,$
- a find the value of the constant $k,$ (3)
- b show that $(x + 1)$ is a factor of $f(x).$ (2)
- 14 $f(x) \equiv x^3 - 4x^2 - 7x + 8.$
- a Find the remainder when $f(x)$ is divided by $(x + 2).$ (2)
- Given that
- $$g(x) \equiv f(x) + c,$$
- and that $(x + 2)$ is a factor of $g(x),$
- b state the value of the constant $c,$ (1)
- c solve the equation $g(x) = 0.$ (4)
- 15 $f(x) \equiv x^3 - 4x + 1.$
- Given that when $f(x)$ is divided by $(2x - k),$ where k is a constant, the remainder is 4,
- a show that $k^3 - 16k - 24 = 0.$ (3)
- Given also that when $f(x)$ is divided by $(x + k)$ the remainder is 1,
- b find the value of $k.$ (3)