

1 a $f'(x) = 6x^2 + 10x$

b $6x^2 + 10x \geq 0$
 $2x(3x + 5) \geq 0$
 $x \leq -\frac{5}{3}$ and $x \geq 0$

3 a $\frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} - 4x^{-2}$

$$\frac{d^2y}{dx^2} = -\frac{1}{4}x^{-\frac{3}{2}} + 8x^{-3}$$

b SP: $\frac{1}{2}x^{-\frac{1}{2}} - 4x^{-2} = 0$
 $\frac{1}{2}x^{-2}(x^{\frac{3}{2}} - 8) = 0$
 $x^{\frac{3}{2}} = 8$
 $x = 4$

$\therefore (4, 3)$

when $x = 4$, $\frac{d^2y}{dx^2} = \frac{3}{32}$

$\frac{d^2y}{dx^2} > 0 \therefore$ minimum

5 a $\frac{dh}{dt} = 8t^3 - 24t^2 + 16t$

b when $t = 0.25$,
 $\frac{dh}{dt} = 2.625$ cm per second

c SP: $8t^3 - 24t^2 + 16t = 0$
 $8t(t - 1)(t - 2) = 0$
 $t = 0, 1, 2$

from graph, max when $t = 1$
 \therefore max height = 3 cm

2 a $\frac{dy}{dx} = 3x^2 - 2x + 2$

at $(1, -2)$, grad = 3

$\therefore y + 2 = 3(x - 1)$

$3x - y - 5 = 0$

b SP when $3x^2 - 2x + 2 = 0$

$b^2 - 4ac = 4 - 24 = -20$

$b^2 - 4ac < 0 \therefore$ no real roots

\therefore no stationary points

4 a $y = 0 \Rightarrow x(x + 3)^2 = 0$

$x = -3, 0$

$\therefore (-3, 0), (0, 0)$


b $f'(x) = 3x^2 + 12x + 9$

decreasing when $3x^2 + 12x + 9 \leq 0$

$3(x + 3)(x + 1) \leq 0$

$\therefore -3 \leq x \leq -1$

c



6 a $\frac{dy}{dx} = 3x^2 + 6kx - 9k^2$

stationary when $3x^2 + 6kx - 9k^2 = 0$

$\Rightarrow x^2 + 2kx - 3k^2 = 0$

b $(x + 3k)(x - k) = 0$

$x = -3k, k$

when $x = k$, $y = k^3 + 3k^3 - 9k^3 = -5k^3$

\therefore stationary at $(k, -5k^3)$

c when $x = -3k$,

$y = -27k^3 + 27k^3 + 27k^3 = 27k^3$

$\therefore (-3k, 27k^3)$

$$7 \quad \mathbf{a} \quad V = \frac{1}{2}x^2 \sin 60^\circ \times l \\ = \frac{1}{2}x^2 l \times \frac{\sqrt{3}}{2} = 250$$

$$\therefore l = \frac{1000}{\sqrt{3}x^2} \text{ or } \frac{1000\sqrt{3}}{3x^2}$$

$$\mathbf{b} \quad A = (2 \times \frac{\sqrt{3}}{4}x^2) + 3xl \\ = \frac{\sqrt{3}}{2}x^2 + (3x \times \frac{1000\sqrt{3}}{3x^2}) \\ = \frac{\sqrt{3}}{2}(x^2 + \frac{2000}{x})$$

$$\mathbf{c} \quad \frac{dA}{dx} = \frac{\sqrt{3}}{2}(2x - 2000x^{-2})$$

$$\text{SP: } \frac{\sqrt{3}}{2}(2x - 2000x^{-2}) = 0$$

$$x^3 = 1000$$

$$x = 10$$

$$\mathbf{d} \quad \min A = 150\sqrt{3}$$

$$\mathbf{e} \quad \frac{d^2A}{dx^2} = \frac{\sqrt{3}}{2}(2 + 4000x^{-3})$$

$$\text{when } x = 10, \frac{d^2A}{dx^2} = 3\sqrt{3}$$

$$\frac{d^2A}{dx^2} > 0 \quad \therefore \text{minimum}$$

$$9 \quad \mathbf{a} \quad x^{\frac{1}{2}} - 4 + 3x^{-\frac{1}{2}} = 0$$

$$x - 4x^{\frac{1}{2}} + 3 = 0$$

$$(x^{\frac{1}{2}} - 1)(x^{\frac{1}{2}} - 3) = 0$$

$$x^{\frac{1}{2}} = 1, 3$$

$$x = 1, 9$$

$$\therefore (1, 0) \text{ and } (9, 0)$$

$$\mathbf{b} \quad \frac{dy}{dx} = \frac{1}{2}x^{-\frac{1}{2}} - \frac{3}{2}x^{-\frac{3}{2}}$$

$$\text{SP: } \frac{1}{2}x^{-\frac{1}{2}} - \frac{3}{2}x^{-\frac{3}{2}} = 0$$

$$\frac{1}{2}x^{-\frac{3}{2}}(x - 3) = 0$$

$$x = 3$$

$$y = \sqrt{3} - 4 + \frac{3}{\sqrt{3}} = 2\sqrt{3} - 4$$

$$\therefore (3, 2\sqrt{3} - 4)$$

$$8 \quad \mathbf{a} \quad f'(x) = 3x^2 + 8x + k$$

for 2 SPs, $f'(x) = 0$ has 2 distinct roots

$$\therefore b^2 - 4ac > 0$$

$$64 - 12k > 0$$

$$k < \frac{16}{3}$$

$$\mathbf{b} \quad \text{SP: } 3x^2 + 8x - 3 = 0$$

$$(3x - 1)(x + 3) = 0$$

$$x = -3, \frac{1}{3}$$

$$\therefore (-3, 19) \text{ and } (\frac{1}{3}, \frac{13}{27})$$

$$10 \quad \mathbf{a} \quad f(-1) = -1 - 3 + 4 = 0$$

$\therefore (x + 1)$ is a factor

$$\mathbf{b} \quad \begin{array}{r} x^2 - 4x + 4 \\ x + 1 \overline{) x^3 - 3x^2 + 0x + 4} \\ \underline{x^3 + x^2} \\ -4x^2 + 0x \\ \underline{-4x^2 - 4x} \\ 4x + 4 \\ \underline{4x + 4} \\ 0 \end{array}$$

$$\therefore f(x) \equiv (x + 1)(x^2 - 4x + 4)$$

$$f(x) \equiv (x + 1)(x - 2)^2$$

$$\mathbf{c} \quad (2, 0), \text{ as } (x - 2) \text{ is a repeated factor}$$

of $f(x)$ so x -axis is a tangent at $(2, 0)$

$$\mathbf{d} \quad f'(x) = 3x^2 - 6x$$

$$\text{SP: } 3x^2 - 6x = 0$$

$$3x(x - 2) = 0$$

$$x = 0, 2$$

$\therefore (0, 4)$ is other turning point