

1 Differentiate with respect to x

- a** $\cos x$ **b** $5 \sin x$ **c** $\cos 3x$ **d** $\sin \frac{1}{4}x$
e $\sin(x+1)$ **f** $\cos(3x-2)$ **g** $4 \sin(\frac{\pi}{3}-x)$ **h** $\cos(\frac{1}{2}x + \frac{\pi}{6})$
i $\sin^2 x$ **j** $2 \cos^3 x$ **k** $\cos^2(x-1)$ **l** $\sin^4 2x$

2 Use the derivatives of $\sin x$ and $\cos x$ to show that

- a** $\frac{d}{dx}(\tan x) = \sec^2 x$ **b** $\frac{d}{dx}(\sec x) = \sec x \tan x$
c $\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$ **d** $\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$

3 Differentiate with respect to t

- a** $\cot 2t$ **b** $\sec(t+2)$ **c** $\tan(4t-3)$ **d** $\operatorname{cosec} 3t$
e $\tan^2 t$ **f** $3 \operatorname{cosec}(t + \frac{\pi}{6})$ **g** $\cot^3 t$ **h** $4 \sec \frac{1}{2}t$
i $\cot(2t-3)$ **j** $\sec^2 2t$ **k** $\frac{1}{2} \tan(\pi-4t)$ **l** $\operatorname{cosec}^2(3t+1)$

4 Differentiate with respect to x

- a** $\ln(\sin x)$ **b** $6e^{\tan x}$ **c** $\sqrt{\cos 2x}$ **d** $e^{\sin 3x}$
e $2 \cot x^2$ **f** $\sqrt{\sec x}$ **g** $3e^{-\operatorname{cosec} 2x}$ **h** $\ln(\tan 4x)$

5 Find the coordinates of any stationary points on each curve in the interval $0 \leq x \leq 2\pi$.

- a** $y = x + 2 \sin x$ **b** $y = 2 \sec x - \tan x$ **c** $y = \sin x + \cos 2x$

6 Find an equation for the tangent to each curve at the point on the curve with the given x -coordinate.

- a** $y = 1 + \sin 2x$, $x = 0$ **b** $y = \cos x$, $x = \frac{\pi}{3}$
c $y = \tan 3x$, $x = \frac{\pi}{4}$ **d** $y = \operatorname{cosec} x - 2 \sin x$, $x = \frac{\pi}{6}$

7 Differentiate with respect to x

- a** $x \sin x$ **b** $\frac{\cos 2x}{x}$ **c** $e^x \cos x$ **d** $\sin x \cos x$
e $x^2 \operatorname{cosec} x$ **f** $\sec x \tan x$ **g** $\frac{x}{\tan x}$ **h** $\frac{\sin 2x}{e^{3x}}$
i $\cos^2 x \cot x$ **j** $\frac{\sec 2x}{x^2}$ **k** $x \tan^2 4x$ **l** $\frac{\sin x}{\cos 2x}$

8 Find the value of $f'(x)$ at the value of x indicated in each case.

- a** $f(x) = \sin 3x \cos 5x$, $x = \frac{\pi}{4}$ **b** $f(x) = \tan 2x \sin x$, $x = \frac{\pi}{3}$
c $f(x) = \frac{\ln(2 \cos x)}{\sin x}$, $x = \frac{\pi}{3}$ **d** $f(x) = \sin^2 x \cos^3 x$, $x = \frac{\pi}{6}$

- 9 Find an equation for the normal to the curve $y = 3 + x \cos 2x$ at the point where it crosses the y -axis.

- 10 A curve has the equation $y = \frac{2 + \sin x}{1 - \sin x}$, $0 \leq x \leq 2\pi$, $x \neq \frac{\pi}{2}$.

- a Find and simplify an expression for $\frac{dy}{dx}$.
- b Find the coordinates of the turning point of the curve.
- c Show that the tangent to the curve at the point P , with x -coordinate $\frac{\pi}{6}$, has equation

$$y = 6\sqrt{3}x + 5 - \sqrt{3}\pi.$$

- 11 A curve has the equation $y = e^{-x} \sin x$.

- a Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$.
- b Find the exact coordinates of the stationary points of the curve in the interval $-\pi \leq x \leq \pi$ and determine their nature.


- 12 The curve C has the equation $y = x \sec x$.

- a Show that the x -coordinate of any stationary point of C must satisfy the equation

$$1 + x \tan x = 0.$$

- b By sketching two suitable graphs on the same set of axes, deduce the number of stationary points C has in the interval $0 \leq x \leq 2\pi$.

13



The diagram shows the curve $y = f(x)$ in the interval $0 \leq x \leq 2\pi$, where

$$f(x) \equiv \cos x \sin 2x.$$

- a Show that $f'(x) = 2 \cos x (1 - 3 \sin^2 x)$.
- b Find the x -coordinates of the stationary points of the curve in the interval $0 \leq x \leq 2\pi$.
- c Show that the maximum value of $f(x)$ in the interval $0 \leq x \leq 2\pi$ is $\frac{4}{9}\sqrt{3}$.
- d Explain why this is the maximum value of $f(x)$ for all real values of x .
- 14 A curve has the equation $y = \operatorname{cosec} \left(x - \frac{\pi}{6}\right)$ and crosses the y -axis at the point P .
- a Find an equation for the normal to the curve at P .
- The point Q on the curve has x -coordinate $\frac{\pi}{3}$.
- b Find an equation for the tangent to the curve at Q .
- The normal to the curve at P and the tangent to the curve at Q intersect at the point R .
- c Show that the x -coordinate of R is given by $\frac{8\sqrt{3} + 4\pi}{13}$.