

- 1 For each equation, show that it can be rearranged into the given iterative form. Use this and the given value of  $x_0$  to find  $x_1$ ,  $x_2$  and  $x_3$ . Give your value of  $x_3$  correct to 4 decimal places.
- a  $9 + 4x - 2x^3 = 0$        $x_{n+1} = \sqrt[3]{2x_n + 4.5}$        $x_0 = 2$
- b  $e^x - 8x + 5 = 0$        $x_{n+1} = \ln(8x_n - 5)$        $x_0 = 3$
- c  $\tan x - 5x + 13 = 0$        $x_{n+1} = \arctan(5x_n - 13)$        $x_0 = -1.2$
- d  $\ln x + \sqrt{x} + 1.4 = 0$        $x_{n+1} = e^{-(\sqrt{x_n} + 1.4)}$        $x_0 = 0.16$
- 2 For each equation, show that it can be rearranged into the given iterative form and state the values of the constants  $a$  and  $b$ . Use this and the given value of  $x_0$  to find  $x_1$ ,  $x_2$  and  $x_3$ . Give your value of  $x_3$  correct to 3 decimal places.
- a  $e^{2x-1} - 6x = 0$        $x_{n+1} = a(\ln bx_n + 1)$        $x_0 = 1.7$
- b  $\frac{2}{x} + \cos x - 3 = 0$        $x_{n+1} = \frac{a}{b - \cos x_n}$        $x_0 = 0.8$
- c  $2x^3 - 6x - 11 = 0$        $x_{n+1} = \sqrt{a + \frac{b}{x_n}}$        $x_0 = 2$
- d  $15 \ln(x + 3) - 4x = 0$        $x_{n+1} = e^{ax_n} + b$        $x_0 = -2.5$
- 3 In each case, use the given iteration formula and value of  $x_0$  to find a root of the equation  $f(x) = 0$  to the stated degree of accuracy. Justify the accuracy of your answers.
- a  $f(x) = 10^x + 3x - 4$        $x_{n+1} = \log_{10}(4 - 3x_n)$        $x_0 = 0.44$       3 decimal places
- b  $f(x) = x^2 + \frac{1}{x-5}$        $x_{n+1} = \sqrt{\frac{x_n^3 + 1}{5}}$        $x_0 = 0.5$       2 significant figures
- c  $f(x) = 30 - 5x + \sin 2x$        $x_{n+1} = 6 + 0.2 \sin 2x_n$        $x_0 = 6$       3 significant figures
- d  $f(x) = e^{4-x} - \ln x$        $x_{n+1} = 4 - \ln(\ln x_n)$        $x_0 = 3.7$       3 decimal places
- 4  $f(x) = x^5 - 10x^3 + 4$ .
- The equation  $f(x) = 0$  has a root in the interval  $-4 < x < -3$ .
- a Use the iteration formula  $x_{n+1} = \sqrt[5]{10x_n^3 - 4}$  and the starting value  $x_0 = -3.2$  to find the value of this root correct to 2 decimal places.
- The equation  $f(x) = 0$  can be rearranged into the iterative form  $x_{n+1} = \sqrt[3]{\frac{a}{b-x_n^2}}$ .
- b Find the values of the constants  $a$  and  $b$  in this formula.
- The equation  $f(x) = 0$  has another root in the interval  $0 < x < 1$ .
- c Using the iteration formula with your values from part **b** and the starting value  $x_0 = 1$ , find the value of this root correct to 3 decimal places.
- 5  $f: x \rightarrow \arcsin 2x - 0.5x - 0.7$ ,  $x \in \mathbb{R}$ ,  $|x| \leq 0.5$
- The equation  $f(x) = 0$  can be rearranged into the iterative form  $x_{n+1} = a \sin(bx_n + c)$ .
- a Find the values of the constants  $a$ ,  $b$  and  $c$  in this formula.
- The equation  $f(x) = 0$  has a solution in the interval  $(0.3, 0.4)$ .
- b Using the iterative formula with your values from part **a** and the starting value  $x_0 = 0.4$ , find this solution correct to 3 decimal places.