


- 1**    **a**  $\frac{dy}{dx} = e^x + 2x$
- b** at A,  $x = 0 \therefore y = -3$ , grad = 1  
 $\therefore y = x - 3$
- c** SP:  $e^x + 2x = 0$   
 let  $f(x) = e^x + 2x$   
 $f(-0.4) = -0.130$   
 $f(-0.3) = 0.141$   
 sign change,  $f(x)$  continuous  $\therefore$  root  
 $\therefore$   $x$ -coord of B in interval  $[-0.4, -0.3]$
- d**  $x_1 = -0.34694$   
 $x_2 = -0.35126$   
 $x_3 = -0.35169$   
 $x_4 = -0.35173$   
 $\therefore$   $x$ -coord of B =  $-0.352$  (3dp)
- 2**    **a**  $f(0) = 0.279$   
 $f(5) = -4.10$   
 $f(1) = 0.266$   
 $f(3) = -2.44$   
 $f(2) = -0.853$   
 $\therefore k = 1$
- b**  $x_0 = 1$   
 $x_1 = 1.2684$   
 $x_2 = 1.3106$   
 $x_3 = 1.3106$
- 3**    **a** area of segment =  $\frac{1}{2}r^2\theta - \frac{1}{2}r^2\sin\theta$   
 $= \frac{1}{2}r^2(\theta - \sin\theta)$   
 $\therefore \frac{1}{2}r^2\sin\theta = 4 \times \frac{1}{2}r^2(\theta - \sin\theta)$   
 $\sin\theta = 4(\theta - \sin\theta)$   
 $\sin\theta = 4\theta - 4\sin\theta$   
 $4\theta - 5\sin\theta = 0$
- b**  $\theta_1 = 1.11401$   
 $\theta_2 = 1.12184$   
 $\theta_3 = 1.12613$   
 $\theta_4 = 1.12844$   
 $\theta_5 = 1.12968$   
 $\therefore \theta = 1.13$  (2dp)
- 4**    **a**  $e^{x^2} - x - 3 = 0$   
 $e^{x^2} = x + 3$   
 $x^2 = \ln(x + 3)$   
 $x = \sqrt{\ln(x + 3)} \therefore a = 1, b = 3$
- b** e.g.  $x_0 = 1.5$   
 $x_1 = 1.226408$   
 $x_2 = 1.200563$   
 $x_3 = 1.198006$   
 $x_4 = 1.197752$   
 $x_5 = 1.197727$   
 $\therefore$  solution =  $1.198$  (3dp)

- 5 a  $y = x^2 - 9$   
 swap  $x = y^2 - 9$   
 $y = \pm\sqrt{x+9}$   
 (domain  $\Rightarrow +$ )  
 $f^{-1}(x) = \sqrt{x+9}$ ,  $x \in \mathbb{R}$ ,  $x \geq -9$   
 range:  $f^{-1}(x) \geq 0$



- c let  $h(x) = f^{-1}(x) + g(x) = \sqrt{x+9} + x^3$   
 $h(-2) = -5.35$   
 $h(-1) = 1.83$   
 sign change,  $h(x)$  continuous  $\therefore$  root  
 d  $x_1 = -1.41421$ ,  $x_2 = -1.40174$ ,  
 $x_3 = -1.40212$ ,  $x_4 = -1.40211$   
 $\therefore$  root =  $-1.402$  (3dp)


- 7 a at A,  $x^{\frac{5}{2}} - 3x^{\frac{1}{2}} - 7x = 0$   
 let  $f(x) = x^{\frac{5}{2}} - 3x^{\frac{1}{2}} - 7x$   
 $f(4) = -2$ ,  $f(5) = 14.2$   
 sign change,  $f(x)$  continuous  $\therefore$  root  
 $\therefore 4 < \alpha < 5$

- b  $\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7$   
 at B,  $\frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7 = 0$   
 let  $g(x) = \frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7$   
 $g(2) = -0.990$ ,  $g(3) = 5.12$   
 sign change,  $g(x)$  continuous  $\therefore$  root  
 $\therefore 2 < \beta < 3$

- c  $\frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7 = 0$   
 $5x^2 - 3 - 14x^{\frac{1}{2}} = 0$   
 $x^2 = 0.6 + 2.8x^{\frac{1}{2}}$   
 $x > 0 \therefore x = \beta$  is a soln to  $x = \sqrt{0.6 + 2.8x^{\frac{1}{2}}}$

- d  $x_1 = 2.158144$   
 $x_2 = 2.171031$   
 $x_3 = 2.173853$   
 $x_4 = 2.174470$   
 $x_5 = 2.174604$   
 $\therefore \beta = 2.175$  (4sf)

- 6 a



- b  $-(-x^2 - 3x) = \frac{1}{x}$   
 $x^2 + 3x = \frac{1}{x}$   
 $x^3 + 3x^2 = 1$   
 $x^3 + 3x^2 - 1 = 0$   
 c  $x_1 = 0.57735$   
 $x_2 = 0.52871$   
 $x_3 = 0.53234$   
 $x_4 = 0.53207$   
 $\therefore$  x-coord of P =  $0.532$  (3dp)

- 8 a  $\frac{dy}{dx} = 3 - \frac{1}{x}$   
 grad = 2  
 $\therefore$  grad of normal =  $-\frac{1}{2}$   
 $\therefore y - 3 = -\frac{1}{2}(x - 1)$   
 $[y = \frac{7}{2} - \frac{1}{2}x]$

- b  $3x - \ln x = \frac{7}{2} - \frac{1}{2}x$

$$6x - 2 \ln x = 7 - x$$

$$2 \ln x - 7x + 7 = 0$$

- c  $2 \ln x = 7x - 7$   
 $\ln x = \frac{7}{2}(x - 1)$   
 $x = e^{\frac{7}{2}(x-1)} \therefore k = \frac{7}{2}$

- d  $x_1 = 0.173774$   
 $x_2 = 0.055477$   
 $x_3 = 0.036669$   
 $x_4 = 0.034333$   
 $x_5 = 0.034053$   
 $\therefore$  x-coord of Q =  $0.034$  (3dp)

- e let  $f(x) = 2 \ln x - 7x + 7$   
 $f(0.0335) = -0.027$   
 $f(0.0345) = 0.025$   
 sign change,  $f(x)$  continuous  $\therefore$  root