

- 1 a e.g.  $a = -2, b = 1 \Rightarrow a^2 - b^2 = 4 - 1 = 3 \Rightarrow a^2 - b^2 > 0$   
 and  $a - b = -2 - 1 = -3 \Rightarrow a - b < 0$   
 [ any negative value of  $a$  such that  $|a| > |b|$  ]
- b 7 7 is prime and divisible by 7 [ no other examples ]
- c e.g.  $x = \sqrt{2}, y = 2\sqrt{2} \Rightarrow x$  and  $y$  irrational  
 and  $xy = 4$  which is rational [ many other examples ]
- d e.g.  $x = -90 \Rightarrow \cos(90 - |x|)^\circ = \cos 0 = 1$   
 and  $\sin x^\circ = \sin(-90^\circ) = -1$  [ any -ve  $x$  except multiples of 180 ]
- 2 a true any number divisible by 6 is also divisible by 2 and  $\therefore$  not prime
- b  $n$  1 2 3 4 5  
 $3^n + 2$  5 11 29 83 245  
 false e.g.  $n = 5 \Rightarrow 3^n + 2 = 245$  which is divisible by 5 and  $\therefore$  not prime  
 [ many other examples ]
- c false e.g.  $n = 4 \Rightarrow \sqrt{n} = 2$  which is rational [ many other examples ]
- d true  $b$  divisible by  $c \Rightarrow b = kc, k \in \mathbb{Z}$   
 $a$  divisible by  $b \Rightarrow a = lb, l \in \mathbb{Z} \Rightarrow a = klc \therefore a$  is divisible by  $c$
- 3 a assume  $n^3$  odd and  $n$  even, where  $n \in \mathbb{Z}^+$   
 $n$  even  $\Rightarrow n = 2m, m \in \mathbb{Z}$   
 $\Rightarrow n^3 = (2m)^3 = 8m^3 = 2(4m^3)$   
 $4m^3 \in \mathbb{Z} \therefore n^3$  even  
 $\Rightarrow$  contradiction  $\therefore n$  odd
- b assume  $x$  irrational and  $\sqrt{x}$  rational  
 $\sqrt{x}$  rational  $\Rightarrow \sqrt{x} = \frac{p}{q}, p, q \in \mathbb{Z}$   
 $\Rightarrow x = \frac{p^2}{q^2}, p^2, q^2 \in \mathbb{Z} \therefore x$  rational  
 $\Rightarrow$  contradiction  $\therefore \sqrt{x}$  irrational
- c assume  $bc$  not divisible by  $a$  and  $b$  divisible by  $a$  where  $a, b, c \in \mathbb{Z}$   
 $b$  divisible by  $a \Rightarrow b = ka, k \in \mathbb{Z}$   
 $\Rightarrow bc = kac$  which is divisible by  $a$   
 $\Rightarrow$  contradiction  $\therefore b$  is not divisible by  $a$
- d assume  $n^2 - 4n$  odd and  $n$  even, where  $n \in \mathbb{Z}^+$   
 $n$  even  $\Rightarrow n = 2m, m \in \mathbb{Z}$   
 $\Rightarrow n^2 - 4n = (2m)^2 - 4(2m) = 4m^2 - 8m = 2(2m^2 - 4m)$   
 $2m^2 - 4m \in \mathbb{Z} \therefore n^2 - 4n$  even  
 $\Rightarrow$  contradiction  $\therefore n$  odd
- e assume  $m^2 - n^2 = 6$ , where  $m, n \in \mathbb{Z}^+$   
 $m^2 - n^2 = 6 \Rightarrow (m+n)(m-n) = 6$   
 $m, n \in \mathbb{Z}^+ \Rightarrow (m+n), (m-n) \in \mathbb{Z}, (m+n) > (m-n)$  and  $(m+n) > 0$   
 $\therefore m+n = 6$  and  $m-n = 1$  or  $m+n = 3$  and  $m-n = 2$   
 adding  $\Rightarrow 2m = 7$  or  $2m = 5$   
 $\Rightarrow m = \frac{7}{2}$  or  $m = \frac{5}{2} \Rightarrow m$  not an integer  
 $\Rightarrow$  contradiction  $\therefore$  no positive integer solutions

- 4 a** assume  $x^2 + y^2$  divisible by 4 and  $x, y$  odd integers  
 $x, y$  odd  $\Rightarrow x = 2m + 1, m \in \mathbb{Z}$  and  $y = 2n + 1, n \in \mathbb{Z}$   
 $\Rightarrow x^2 + y^2 = (2m + 1)^2 + (2n + 1)^2$   
 $= 4m^2 + 4m + 1 + 4n^2 + 4n + 1$   
 $= 4(m^2 + m + n^2 + n) + 2$   
 $m^2 + m + n^2 + n \in \mathbb{Z} \therefore x^2 + y^2$  not divisible by 4  
 $\Rightarrow$  contradiction  $\therefore x$  and  $y$  not both odd
- b** assume  $x^2 + y^2$  divisible by 4,  $x$  odd integer and  $y$  even integer  
 $x$  odd,  $y$  even  $\Rightarrow x = 2m + 1, m \in \mathbb{Z}$  and  $y = 2n, n \in \mathbb{Z}$   
 $\Rightarrow x^2 + y^2 = (2m + 1)^2 + (2n)^2$   
 $= 4m^2 + 4m + 1 + 4n^2$   
 $= 4(m^2 + m + n^2) + 1$   
 $m^2 + m + n^2 \in \mathbb{Z} \therefore x^2 + y^2$  not divisible by 4  
 $\Rightarrow$  contradiction  $\therefore x$  odd and  $y$  even not possible  
 same argument applies with  $x$  even and  $y$  odd  
 part **a** shows  $x$  and  $y$  can't both be odd  
 $\therefore x$  and  $y$  both even
- 5 a** false e.g.  $a = 2, b = 4 \Rightarrow \log_a b = 2$  which is rational  
 [ many other examples ]
- b** true  $(2n + 1)$  and  $(2n + 3), n \in \mathbb{Z}$  represent any two consecutive odd integers  
 $(2n + 3)^2 - (2n + 1)^2 = 4n^2 + 12n + 9 - (4n^2 + 4n + 1)$   
 $= 8n + 8$   
 $= 8(n + 1)$   
 $n + 1 \in \mathbb{Z} \therefore$  difference is divisible by 8
- c** false e.g.  $n = 13 \Rightarrow n^2 + 3n + 13 = 13(13 + 3 + 1)$  which is divisible by 13  
 [ many other examples ]
- d** true  $x^2 - 2y(x - y) = x^2 - 2xy + 2y^2$   
 $= x^2 - 2xy + y^2 + y^2$   
 $= (x - y)^2 + y^2$   
 for real  $x$  and  $y, (x - y)^2 \geq 0$  and  $y^2 \geq 0 \therefore x^2 - 2y(x - y) \geq 0$
- 6 a**  $\sqrt{2} = \frac{p}{q}, p, q \in \mathbb{Z} \Rightarrow 2 = \frac{p^2}{q^2} \Rightarrow p^2 = 2q^2$   
 $\Rightarrow p^2$  even  $\Rightarrow p$  even
- b** assume  $\sqrt{2}$  rational  $\Rightarrow \sqrt{2} = \frac{p}{q}, p, q \in \mathbb{Z}$  and  $p, q$  co-prime  
 part **a**  $\Rightarrow p$  even  $\Rightarrow p = 2n, n \in \mathbb{Z}$   
 $\Rightarrow (2n)^2 = 2q^2$   
 $\Rightarrow q^2 = 2n^2$   
 $\Rightarrow q^2$  even  $\Rightarrow q$  even  
 $\Rightarrow p$  and  $q$  both even  $\therefore$  not co-prime  
 $\Rightarrow$  contradiction  $\therefore \sqrt{2}$  is irrational