

$$1 \quad \mathbf{a} \quad \frac{3x+5}{(x+1)(x+3)} \equiv \frac{A}{x+1} + \frac{B}{x+3}$$

$$3x+5 \equiv A(x+3) + B(x+1)$$

$$x = -1 \Rightarrow 2 = 2A \Rightarrow A = 1$$

$$x = -3 \Rightarrow -4 = -2B \Rightarrow B = 2$$

$$\therefore \frac{3x+5}{(x+1)(x+3)} \equiv \frac{1}{x+1} + \frac{2}{x+3}$$

$$\mathbf{b} = \int \left(\frac{1}{x+1} + \frac{2}{x+3} \right) dx$$

$$= \ln|x+1| + 2 \ln|x+3| + c$$

$$2 \quad \frac{3}{(t-2)(t+1)} \equiv \frac{A}{t-2} + \frac{B}{t+1}$$

$$3 \equiv A(t+1) + B(t-2)$$

$$t = 2 \Rightarrow 3 = 3A \Rightarrow A = 1$$

$$t = -1 \Rightarrow 3 = -3B \Rightarrow B = -1$$

$$\therefore \int \frac{3}{(t-2)(t+1)} dt$$

$$= \int \left(\frac{1}{t-2} - \frac{1}{t+1} \right) dt$$

$$= \ln|t-2| - \ln|t+1| + c$$

$$= \ln \left| \frac{t-2}{t+1} \right| + c$$

$$3 \quad \mathbf{a} \quad \frac{6x-11}{(2x+1)(x-3)} \equiv \frac{A}{2x+1} + \frac{B}{x-3}$$

$$6x-11 \equiv A(x-3) + B(2x+1)$$

$$x = -\frac{1}{2} \Rightarrow -14 = -\frac{7}{2}A \Rightarrow A = 4$$

$$x = 3 \Rightarrow 7 = 7B \Rightarrow B = 1$$

$$\therefore \int \frac{6x-11}{(2x+1)(x-3)} dx$$

$$= \int \left(\frac{4}{2x+1} + \frac{1}{x-3} \right) dx$$

$$= 2 \ln|2x+1| + \ln|x-3| + c$$

$$\mathbf{b} \quad \frac{14-x}{x^2+2x-8} \equiv \frac{A}{x+4} + \frac{B}{x-2}$$

$$14-x \equiv A(x-2) + B(x+4)$$

$$x = -4 \Rightarrow 18 = -6A \Rightarrow A = -3$$

$$x = 2 \Rightarrow 12 = 6B \Rightarrow B = 2$$

$$\therefore \int \frac{14-x}{x^2+2x-8} dx$$

$$= \int \left(\frac{2}{x-2} - \frac{3}{x+4} \right) dx$$

$$= 2 \ln|x-2| - 3 \ln|x+4| + c$$

$$\mathbf{c} \quad \frac{6}{(2+x)(1-x)} \equiv \frac{A}{2+x} + \frac{B}{1-x}$$

$$6 \equiv A(1-x) + B(2+x)$$

$$x = -2 \Rightarrow 6 = 3A \Rightarrow A = 2$$

$$x = 1 \Rightarrow 6 = 3B \Rightarrow B = 2$$

$$\therefore \int \frac{6}{(2+x)(1-x)} dx$$

$$= \int \left(\frac{2}{2+x} + \frac{2}{1-x} \right) dx$$

$$= 2 \ln|2+x| - 2 \ln|1-x| + c$$

$$= 2 \ln \left| \frac{2+x}{1-x} \right| + c$$

$$\mathbf{d} \quad \frac{x+1}{5x^2-14x+8} \equiv \frac{A}{5x-4} + \frac{B}{x-2}$$

$$x+1 \equiv A(x-2) + B(5x-4)$$

$$x = \frac{4}{5} \Rightarrow \frac{9}{5} = -\frac{6}{5}A \Rightarrow A = -\frac{3}{2}$$

$$x = 2 \Rightarrow 3 = 6B \Rightarrow B = \frac{1}{2}$$

$$\therefore \int \frac{x+1}{5x^2-14x+8} dx$$

$$= \int \left(\frac{\frac{1}{2}}{(x-2)} - \frac{\frac{3}{2}}{5x-4} \right) dx$$

$$= \frac{1}{2} \ln|x-2| - \frac{3}{10} \ln|5x-4| + c$$

$$4 \quad \mathbf{a} \quad x^2 - 6 \equiv A(x+4)(x-1) + B(x-1) + C(x+4)$$

$$x = -4 \Rightarrow 10 = -5B \Rightarrow B = -2$$

$$x = 1 \Rightarrow -5 = 5C \Rightarrow C = -1$$

$$\text{coeffs of } x^2 \Rightarrow A = 1$$

$$\mathbf{b} = \int \left(1 - \frac{2}{x+4} - \frac{1}{x-1} \right) dx$$

$$= x - 2 \ln|x+4| - \ln|x-1| + c$$

5 a $\frac{x^2-x-4}{(x+2)(x+1)^2} \equiv \frac{A}{x+2} + \frac{B}{x+1} + \frac{C}{(x+1)^2}$
 $x^2-x-4 \equiv A(x+1)^2 + B(x+2)(x+1) + C(x+2)$
 $x = -2 \Rightarrow A = 2$
 $x = -1 \Rightarrow C = -2$
 coeffs of $x^2 \Rightarrow 1 = A + B \Rightarrow B = -1$
 $\therefore \frac{x^2-x-4}{(x+2)(x+1)^2} \equiv \frac{2}{x+2} - \frac{1}{x+1} - \frac{2}{(x+1)^2}$

b $= \int \left(\frac{2}{x+2} - \frac{1}{x+1} - \frac{2}{(x+1)^2} \right) dx$
 $= 2 \ln|x+2| - \ln|x+1| + 2(x+1)^{-1} + c$

6 a $\frac{3x^2-5}{x^2-1} \equiv A + \frac{B}{x+1} + \frac{C}{x-1}$
 $3x^2-5 \equiv A(x+1)(x-1) + B(x-1) + C(x+1)$
 $x = -1 \Rightarrow -2 = -2B \Rightarrow B = 1$
 $x = 1 \Rightarrow -2 = 2C \Rightarrow C = -1$
 coeffs of $x^2 \Rightarrow A = 3$

$\therefore \int \frac{3x^2-5}{x^2-1} dx = \int \left(3 + \frac{1}{x+1} - \frac{1}{x-1} \right) dx$
 $= 3x + \ln|x+1| - \ln|x-1| + c = 3x + \ln \left| \frac{x+1}{x-1} \right| + c$

b $\frac{x(4x+13)}{(2+x)^2(3-x)} \equiv \frac{A}{2+x} + \frac{B}{(2+x)^2} + \frac{C}{3-x}$
 $x(4x+13) \equiv A(2+x)(3-x) + B(3-x) + C(2+x)^2$
 $x = -2 \Rightarrow -10 = 5B \Rightarrow B = -2$
 $x = 3 \Rightarrow 75 = 25C \Rightarrow C = 3$
 coeffs of $x^2 \Rightarrow 4 = -A + C \Rightarrow A = -1$
 $\therefore \int \frac{x(4x+13)}{(2+x)^2(3-x)} dx = \int \left(\frac{3}{3-x} - \frac{1}{2+x} - \frac{2}{(2+x)^2} \right) dx$
 $= -3 \ln|3-x| - \ln|2+x| + 2(2+x)^{-1} + c$

c $\frac{x^2-x+1}{x^2-3x-10} \equiv A + \frac{B}{x-5} + \frac{C}{x+2}$
 $x^2-x+1 \equiv A(x-5)(x+2) + B(x+2) + C(x-5)$
 $x = 5 \Rightarrow 21 = 7B \Rightarrow B = 3$
 $x = -2 \Rightarrow 7 = -7C \Rightarrow C = -1$
 coeffs of $x^2 \Rightarrow A = 1$
 $\therefore \int \frac{x^2-x+1}{x^2-3x-10} dx = \int \left(1 + \frac{3}{x-5} - \frac{1}{x+2} \right) dx$
 $= x + 3 \ln|x-5| - \ln|x+2| + c$

d $\frac{2-6x+5x^2}{x^2(1-2x)} \equiv \frac{A}{x} + \frac{B}{x^2} + \frac{C}{1-2x}$
 $2-6x+5x^2 \equiv Ax(1-2x) + B(1-2x) + Cx^2$
 $x = \frac{1}{2} \Rightarrow \frac{1}{4} = \frac{1}{4}C \Rightarrow C = 1$
 $x = 0 \Rightarrow B = 2$
 coeffs of $x^2 \Rightarrow 5 = -2A + C \Rightarrow A = -2$
 $\therefore \int \frac{2-6x+5x^2}{x^2(1-2x)} dx = \int \left(\frac{1}{1-2x} - \frac{2}{x} + \frac{2}{x^2} \right) dx$
 $= -\frac{1}{2} \ln|1-2x| - 2 \ln|x| - 2x^{-1} + c$

$$7 \quad \frac{3x-5}{(x-1)(x-2)} \equiv \frac{A}{x-1} + \frac{B}{x-2}$$

$$3x-5 \equiv A(x-2) + B(x-1)$$

$$x=1 \quad \Rightarrow \quad -2 = -A \quad \Rightarrow \quad A=2$$

$$x=2 \quad \Rightarrow \quad -1 = B \quad \Rightarrow \quad B=1$$

$$\therefore \int_3^4 \frac{3x-5}{(x-1)(x-2)} dx = \int_3^4 \left(\frac{2}{x-1} + \frac{1}{x-2} \right) dx$$

$$= [2 \ln|x-1| + \ln|x-2|]_3^4$$

$$= (2 \ln 3 + \ln 2) - (2 \ln 2 + 0) = 2 \ln 3 - \ln 2$$

$$8 \quad \text{a} \quad \frac{x+3}{x(x+1)} \equiv \frac{A}{x} + \frac{B}{x+1}$$

$$x+3 \equiv A(x+1) + Bx$$

$$x=0 \quad \Rightarrow \quad 3 = A \quad \Rightarrow \quad A=3$$

$$x=-1 \quad \Rightarrow \quad 2 = -B \quad \Rightarrow \quad B=-2$$

$$\therefore \int_1^3 \frac{x+3}{x(x+1)} dx = \int_1^3 \left(\frac{3}{x} - \frac{2}{x+1} \right) dx$$

$$= [3 \ln|x| - 2 \ln|x+1|]_1^3$$

$$= (3 \ln 3 - 2 \ln 4) - (0 - 2 \ln 2) = 3 \ln 3 - 2 \ln 2$$

$$\text{b} \quad \frac{3x-2}{x^2+x-12} \equiv \frac{A}{x+4} + \frac{B}{x-3}$$

$$3x-2 \equiv A(x-3) + B(x+4)$$

$$x=-4 \quad \Rightarrow \quad -14 = -7A \quad \Rightarrow \quad A=2$$

$$x=3 \quad \Rightarrow \quad 7 = 7B \quad \Rightarrow \quad B=1$$

$$\therefore \int_0^2 \frac{3x-2}{x^2+x-12} dx = \int_0^2 \left(\frac{2}{x+4} + \frac{1}{x-3} \right) dx$$

$$= [2 \ln|x+4| + \ln|x-3|]_0^2$$

$$= (2 \ln 6 + 0) - (2 \ln 4 + \ln 3)$$

$$= 2(\ln 2 + \ln 3) - 4 \ln 2 - \ln 3 = \ln 3 - 2 \ln 2$$

$$\text{c} \quad \frac{9}{2x^2-7x-4} \equiv \frac{A}{2x+1} + \frac{B}{x-4}$$

$$9 \equiv A(x-4) + B(2x+1)$$

$$x = -\frac{1}{2} \quad \Rightarrow \quad 9 = -\frac{9}{2}A \quad \Rightarrow \quad A = -2$$

$$x=4 \quad \Rightarrow \quad 9 = 9B \quad \Rightarrow \quad B=1$$

$$\therefore \int_1^2 \frac{9}{2x^2-7x-4} dx = \int_1^2 \left(\frac{1}{x-4} - \frac{2}{2x+1} \right) dx$$

$$= [\ln|x-4| - \ln|2x+1|]_1^2$$

$$= (\ln 2 - \ln 5) - (\ln 3 - \ln 3) = \ln 2 - \ln 5$$

$$\text{d} \quad \frac{2x^2-7x+7}{x^2-2x-3} \equiv A + \frac{B}{x-3} + \frac{C}{x+1}$$

$$2x^2-7x+7 \equiv A(x-3)(x+1) + B(x+1) + C(x-3)$$

$$x=3 \quad \Rightarrow \quad 4 = 4B \quad \Rightarrow \quad B=1$$

$$x=-1 \quad \Rightarrow \quad 16 = -4C \quad \Rightarrow \quad C=-4$$

$$\text{coeffs of } x^2 \quad \Rightarrow \quad A=2$$

$$\therefore \int_0^2 \frac{2x^2-7x+7}{x^2-2x-3} dx = \int_0^2 \left(2 + \frac{1}{x-3} - \frac{4}{x+1} \right) dx$$

$$= [2x + \ln|x-3| - 4 \ln|x+1|]_0^2$$

$$= (4 + 0 - 4 \ln 3) - (0 + \ln 3 - 0) = 4 - 5 \ln 3$$

$$\begin{aligned} \text{e } \frac{5x+7}{(x+1)^2(x+3)} &\equiv \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{x+3} \\ 5x+7 &\equiv A(x+1)(x+3) + B(x+3) + C(x+1)^2 \\ x=-1 &\Rightarrow 2=2B \Rightarrow B=1 \\ x=-3 &\Rightarrow -8=4C \Rightarrow C=-2 \\ \text{coeffs of } x^2 &\Rightarrow 0=A+C \Rightarrow A=2 \\ \therefore \int_0^1 \frac{5x+7}{(x+1)^2(x+3)} dx &= \int_0^1 \left(\frac{2}{x+1} + \frac{1}{(x+1)^2} - \frac{2}{x+3} \right) dx \\ &= [2 \ln|x+1| - (x+1)^{-1} - 2 \ln|x+3|]_0^1 \\ &= (2 \ln 2 - \frac{1}{2} - 2 \ln 4) - (0 - 1 - 2 \ln 3) \\ &= \frac{1}{2} - 2 \ln 2 + 2 \ln 3 \end{aligned}$$

$$\begin{aligned} \text{f } \frac{2+x}{8-2x-x^2} &\equiv \frac{A}{4+x} + \frac{B}{2-x} \\ 2+x &\equiv A(2-x) + B(4+x) \\ x=-4 &\Rightarrow -2=6A \Rightarrow A=-\frac{1}{3} \\ x=2 &\Rightarrow 4=6B \Rightarrow B=\frac{2}{3} \\ \therefore \int_{-1}^1 \frac{2+x}{8-2x-x^2} dx &= \int_{-1}^1 \left(\frac{\frac{2}{3}}{2-x} - \frac{\frac{1}{3}}{4+x} \right) dx \\ &= \left[-\frac{2}{3} \ln|2-x| - \frac{1}{3} \ln|4+x| \right]_{-1}^1 \\ &= (0 - \frac{1}{3} \ln 5) - (-\frac{2}{3} \ln 3 - \frac{1}{3} \ln 3) \\ &= \ln 3 - \frac{1}{3} \ln 5 \end{aligned}$$

$$\begin{aligned} \text{9 a } \frac{1}{x^2-a^2} &\equiv \frac{A}{x+a} + \frac{B}{x-a} \\ 1 &\equiv A(x-a) + B(x+a) \\ x=-a &\Rightarrow 1=-2aA \Rightarrow A=-\frac{1}{2a} \\ x=a &\Rightarrow 1=2aB \Rightarrow B=\frac{1}{2a} \\ \therefore \frac{1}{x^2-a^2} &\equiv \frac{1}{2a(x-a)} - \frac{1}{2a(x+a)} \end{aligned}$$

$$\begin{aligned} \text{b } \int \frac{1}{x^2-a^2} dx &= \frac{1}{2a} \int \left(\frac{1}{x-a} - \frac{1}{x+a} \right) dx \\ &= \frac{1}{2a} (\ln|x-a| - \ln|x+a|) + c \\ &= \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c \end{aligned}$$

$$\begin{aligned} \text{c } \int \frac{1}{a^2-x^2} dx &= -\int \frac{1}{x^2-a^2} dx \\ &= -\frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c \\ &= \frac{1}{2a} \ln \left| \frac{x+a}{x-a} \right| + c \end{aligned}$$

$$\begin{aligned} \text{10 a } &= \left[\frac{1}{6} \ln \left| \frac{x-3}{x+3} \right| \right]_{-1}^1 \\ &= \frac{1}{6} (\ln \frac{1}{2} - \ln 2) \\ &= -\frac{1}{3} \ln 2 \end{aligned}$$

$$\begin{aligned} \text{b } &= 4 \left[\frac{1}{2} \ln \left| \frac{x+1}{x-1} \right| \right]_{-\frac{1}{2}}^{\frac{1}{2}} \\ &= 2 (\ln 3 - \ln \frac{1}{3}) \\ &= 4 \ln 3 \end{aligned}$$

$$\begin{aligned} \text{c } &= \frac{3}{2} \left[\frac{1}{4} \ln \left| \frac{x-2}{x+2} \right| \right]_0^1 \\ &= \frac{3}{8} (\ln \frac{1}{3} - 0) \\ &= -\frac{3}{8} \ln 3 \end{aligned}$$