

- 1 a Expand $(1 - 4x)^{\frac{1}{2}}$ in ascending powers of x up to and including the term in x^3 and state the set of values of x for which the expansion is valid. (4)

- b By substituting $x = 0.01$ in your expansion, find the value of $\sqrt{6}$ to 6 significant figures. (3)

2
$$f(x) \equiv \frac{4}{1 + 2x - 3x^2}.$$

- a Express $f(x)$ in partial fractions. (3)

- b Hence, or otherwise, find the series expansion of $f(x)$ in ascending powers of x up to and including the term in x^3 and state the set of values of x for which the expansion is valid. (5)

- 3 a Expand $(2 - x)^{-2}$, $|x| < 2$, in ascending powers of x up to and including the term in x^3 . (4)

- b Hence, find the coefficient of x^3 in the series expansion of $\frac{3 - x}{(2 - x)^2}$. (2)

4
$$f(x) \equiv \frac{4}{\sqrt{1 + \frac{2}{3}x}}, \quad -\frac{3}{2} < x < \frac{3}{2}.$$

- a Show that $f(\frac{1}{10}) = \sqrt{15}$. (2)

- b Expand $f(x)$ in ascending powers of x up to and including the term in x^2 . (3)

- c Use your expansion to obtain an approximation for $\sqrt{15}$, giving your answer as an exact, simplified fraction. (2)

- d Show that $3\frac{55}{63}$ is a more accurate approximation for $\sqrt{15}$. (2)

- 5 a Expand $(1 - x)^{\frac{1}{3}}$, $|x| < 1$, in ascending powers of x up to and including the term in x^2 . (3)

- b By substituting $x = 10^{-3}$ in your expansion, find the cube root of 37 correct to 9 significant figures. (3)

- 6 The series expansion of $(1 + 5x)^{\frac{3}{5}}$, in ascending powers of x up to and including the term in x^3 , is

$$1 + 3x + px^2 + qx^3, \quad |x| < \frac{1}{5}.$$

- a Find the values of the constants p and q . (4)

- b Use the expansion with a suitable value of x to find an approximate value for $(1.1)^{\frac{3}{5}}$. (2)

- c Obtain the value of $(1.1)^{\frac{3}{5}}$ from your calculator and hence find the percentage error in your answer to part b. (2)

- 7 a Find the values of A , B and C such that

$$\frac{8 - 6x^2}{(1 + x)(2 + x)^2} \equiv \frac{A}{1 + x} + \frac{B}{2 + x} + \frac{C}{(2 + x)^2}. \quad (4)$$

- b Hence find the series expansion of $\frac{8 - 6x^2}{(1 + x)(2 + x)^2}$, $|x| < 1$, in ascending powers of x up to and including the term in x^3 , simplifying each coefficient. (7)

- 8 a Expand $(1 - 2x)^{\frac{1}{2}}$, $|x| < \frac{1}{2}$, in ascending powers of x up to and including the term in x^2 . (3)
 b By substituting $x = 0.0008$ in your expansion, find the square root of 39 correct to 7 significant figures. (4)

- 9 a Find the series expansion of $(1 + 8x)^{\frac{1}{3}}$, $|x| < \frac{1}{8}$, in ascending powers of x up to and including the term in x^2 , simplifying each term. (3)
 b Find the exact fraction k such that

$$\sqrt[3]{5} = k\sqrt[3]{1.08} \quad (2)$$

- c Hence, use your answer to part **a** together with a suitable value of x to obtain an estimate for $\sqrt[3]{5}$, giving your answer to 4 significant figures. (3)

10
$$f(x) \equiv \frac{6x}{x^2 - 4x + 3}, \quad |x| < 1.$$

- a Express $f(x)$ in partial fractions. (3)
 b Show that for small values of x ,

$$f(x) \approx 2x + \frac{8}{3}x^2 + \frac{26}{9}x^3. \quad (5)$$

- 11 a Find the binomial expansion of $(4 + x)^{\frac{1}{2}}$ in ascending powers of x up to and including the term in x^2 and state the set of values of x for which the expansion is valid. (4)
 b By substituting $x = \frac{1}{20}$ in your expansion, find an estimate for $\sqrt{5}$, giving your answer to 9 significant figures. (3)
 c Obtain the value of $\sqrt{5}$ from your calculator and hence comment on the accuracy of the estimate found in part **b**. (2)

- 12 a Expand $(1 + 2x)^{-\frac{1}{2}}$, $|x| < \frac{1}{2}$, in ascending powers of x up to and including the term in x^3 . (4)
 b Hence, show that for small values of x ,

$$\frac{2 - 5x}{\sqrt{1 + 2x}} \approx 2 - 7x + 8x^2 - \frac{25}{2}x^3. \quad (3)$$

- c Solve the equation

$$\frac{2 - 5x}{\sqrt{1 + 2x}} = \sqrt{3}. \quad (3)$$

- d Use your answers to parts **b** and **c** to find an approximate value for $\sqrt{3}$. (2)

- 13 a Expand $(1 + x)^{-1}$, $|x| < 1$, in ascending powers of x up to and including the term in x^3 . (2)
 b Hence, write down the first four terms in the expansion in ascending powers of x of $(1 + bx)^{-1}$, where b is a constant, for $|bx| < 1$. (1)

Given that in the series expansion of

$$\frac{1 + ax}{1 + bx}, \quad |bx| < 1,$$

the coefficient of x is -4 and the coefficient of x^2 is 12 ,

- c find the values of the constants a and b , (5)
 d find the coefficient of x^3 in the expansion. (2)