

Name:

Date:

C5 - Test 6  
ENERGY CHANGES  
Advanced

**GCSE**

**CHEMISTRY**

**AQA - Triple Science**

Mark

Grade

---

### Materials

For this paper you must have:

- Ruler
- Pencil and Rubber
- Scientific calculator, which you are expected to use when appropriate

### Instructions


- Answer all questions
- Answer questions in the space provided
- All working must be shown

### Information

- The marks for the questions are shown in brackets

**1.**

A student investigated simple cells using the apparatus shown in the figure below.




- If metal 2 is more reactive than metal 1 then the voltage measured is positive.
- If metal 1 is more reactive than metal 2 then the voltage measured is negative.
- The bigger the difference in reactivity of the two metals, the larger the voltage produced.

The student's results are shown in the table below.

Metal 1 \ Metal 2	Chromium	Copper	Iron	Tin	Zinc
Chromium	0.0 V				
Copper	1.2 V	0.0 V			
Iron	0.5 V	not measured	0.0 V		
Tin	0.8 V	-0.4 V	0.3 V	0.0 V	
Zinc	0.2 V	-1.0 V	-0.3 V	-0.6 V	0.0 V

- (a) The ionic equation for the reaction occurring at the zinc electrode in the simple cell made using copper and zinc electrodes is:



Zinc is oxidised in this reaction.

Give a reason why this is oxidation.

---

(1)

(b) Look at the table above.

Which **one** of the metals used was the least reactive?

Give a reason for your answer.

Metal \_\_\_\_\_

Reason \_\_\_\_\_

---

---

(2)

(c) Predict the voltage that would be obtained for a simple cell that has iron as metal **1** and copper as metal **2**.

Explain your answer.

---

---

---

---

---

---

---

---

(3)

(d) Hydrogen fuel cells have been developed for cars.

Write a word equation for the overall reaction that takes place in a hydrogen fuel cell.

---

(1)

- (e) Write the **two** half equations for the reactions that occur at the electrodes in a hydrogen fuel cell.

---

---

---

---

(2)

(Total 9 marks)

2.

Some students investigated the energy changes occurring in the reaction between potassium hydrogencarbonate and hydrochloric acid.

The equation for the reaction is:



This is the method used.

1. Measure 50 cm<sup>3</sup> hydrochloric acid into a glass beaker.
2. Measure the temperature of the hydrochloric acid.
3. Measure a given mass of potassium hydrogencarbonate.
4. Add the potassium hydrogencarbonate to the hydrochloric acid.
5. Stir until all the potassium hydrogencarbonate has reacted.
6. Record the lowest temperature reached.
7. Repeat three more times, using the same mass of potassium hydrogencarbonate.

Each student used a different mass of potassium hydrogencarbonate.

- (a) The method described will not give very accurate results.

Suggest **one** change to the apparatus that would improve the accuracy of the results.

Give a reason for your answer.

---

---

---

---

(2)

(b) The students controlled the volume of the hydrochloric acid.

Give **one** other control variable the students should use.

---

---

(1)

(c) The table shows one student's results.

	<b>Trial 1</b>	<b>Trial 2</b>	<b>Trial 3</b>	<b>Trial 4</b>
<b>Initial temperature in °C</b>	21.2	21.1	21.0	21.1
<b>Final temperature in °C</b>	15.6	15.4	15.6	16.6
<b>Temperature decrease in °C</b>	5.6	5.7	5.4	4.5

Calculate the mean temperature decrease for the results shown in the table above.

Ignore any anomalous results.

Give your answer to 1 decimal place.

Give the uncertainty in your answer.

---


---

---

Mean = \_\_\_\_\_ °C ± \_\_\_\_\_ °C

(3)

The graph below shows the students' results.



(d) Draw **two** intersecting straight lines of best fit on the graph above.

(2)

(e) Explain why the graph has this shape.

Use data from the graph.

---

---

---

---

---

---

---

(3)

(f) Suggest a possible reason for the anomalous points.

Do **not** include errors in measuring.

---


---

(1)


(Total 12 marks)

**3.** This question is about ethanol.

(a) Ethanol is produced by the reaction of ethene and steam:



(i) **Figure 1** shows the energy level diagram for the reaction.



How does the energy level diagram show that the reaction is exothermic?

---

---

(1)

(ii) A catalyst is used for the reaction.

Explain how a catalyst increases the rate of the reaction.

---


---

---

---


(2)

(b) **Figure 2** shows the displayed structure of ethanol.



Complete the dot and cross diagram in **Figure 3** to show the bonding in ethanol.

Show the outer shell electrons only.




(2)



(c) A student burned some ethanol.

**Figure 4** shows the apparatus the student used.



- (i) The student recorded the temperature of the water before and after heating. His results are shown in **Table 1**.

**Table 1**

Temperature before heating	20.7 °C
Temperature after heating	35.1 °C

Calculate the energy used to heat the water.

Use the equation  $Q = m \times c \times \Delta T$

The specific heat capacity of water = 4.2 J / g / °C

---

---

---

---

---

Energy used = \_\_\_\_\_ J

**(3)**

- (ii) **Table 2** shows the mass of the spirit burner before the ethanol was burned and after the ethanol was burned.

**Table 2**

Mass of spirit burner before ethanol was burned	72.80 g
Mass of spirit burner after ethanol was burned	72.10 g

Calculate the number of moles of ethanol ( $C_2H_5OH$ ) that were burned.

Relative atomic masses ( $A_r$ ): H = 1; C = 12; O = 16

---

---

---

---

---

Number of moles burned = \_\_\_\_\_

**(3)**

- (iii) Calculate the energy released in joules per mole.

You should assume that all the energy from the ethanol burning was used to heat the water.

---

Energy = \_\_\_\_\_ J / mole

**(1)**

- (d) The names, structures and boiling points of ethanol and two other alcohols are shown in **Table 3**.

**Table 3**

Name	Methanol	Ethanol	Propanol
Structure	$\begin{array}{c} \text{H} \\   \\ \text{H}-\text{C}-\text{O}-\text{H} \\   \\ \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\   \quad   \\ \text{H} \quad \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{O}-\text{H} \\   \quad   \quad   \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$
Boiling point in °C	65	78	97

Use your knowledge of structure and bonding to suggest why the boiling points increase as the number of carbon atoms increases.

---

---

---

---

---

---

---

---


(3)

(Total 15 marks)

4.

This question is about energy changes in chemical reactions.


- (a) Balance the chemical equation for the combustion of methane.



(1)

(b) Alcohols are used as fuels.

A group of students investigated the amount of energy released when an alcohol was burned. The students used the apparatus shown in the diagram below.



In one experiment the temperature of 50 g of water increased from 22.0 °C to 38.4 °C. The mass of alcohol burned was 0.8 g.

Calculate the heat energy (Q) in joules, released by burning 0.8 g of the alcohol. Use the equation:

$$Q = m \times c \times \Delta T$$

Specific heat capacity (c) = 4.2 J / g / °C

---

---


---

---


Heat energy (Q) = \_\_\_\_\_ J

(3)

(c) The chemical equation for the combustion of ethanol is:



(i) The equation for the reaction can be shown as:



Bond	Bond energy in kJ per mole
C — H	413
C — C	347
C — O	358
C = O	799
O — H	467
O = O	495

Use the bond energies to calculate the overall energy change for this reaction.

---

---

---

---

---

---

---

Overall energy change = \_\_\_\_\_ kJ per mole

(3)

(ii) The reaction is exothermic.  
Explain why, in terms of bonds broken and bonds formed.

---

---

---


---

(2)

(iii) Complete the energy level diagram for the combustion of ethanol.

On the completed diagram, label:


- activation energy
- overall energy change.




(3)  
(Total 12 marks)

**5.**

A Bunsen burner releases heat energy by burning methane in air.



- (a) Methane (CH<sub>4</sub>) reacts with oxygen from the air to produce carbon dioxide and water.
- (i) Use the equation and the bond energies to calculate a value for the energy change in this reaction.



Bond	Bond energy in kJ per mole
C — H	414
O = O	498
C = O	803
O—H	464

---

---

---

---

---

---

---

---

---

Energy change = \_\_\_\_\_ kJ per mole

**(3)**

(ii) This reaction releases heat energy.

Explain why, in terms of bond energies.

---

---

---

---

**(2)**

(b) If the gas tap to the Bunsen burner is turned on, the methane does not start burning until it is lit with a match.

Why is heat from the match needed to start the methane burning?

---

---

**(1)**


**(Total 6 marks)**



6.

This question is about the reaction of ethene and bromine.

The equation for the reaction is:




(a) Complete the reaction profile in **Figure 1**.

Draw labelled arrows to show:

- The energy given out ( $\Delta H$ )
- The activation energy.

**Figure 1**



(3)

(b) When ethene reacts with bromine, energy is required to break covalent bonds in the molecules.

Explain how a covalent bond holds two atoms together.

---

---


---

---

(2)

- (c) **Figure 2** shows the displayed formulae for the reaction of ethene with bromine.

**Figure 2**



The bond enthalpies and the overall energy change are shown in the table below.

	<b>C=C</b>	<b>C-H</b>	<b>C-C</b>	<b>C-Br</b>	<b>Overall energy change</b>
<b>Energy in kJ / mole</b>	612	412	348	276	-95

Use the information in the table above and **Figure 2** to calculate the bond energy for the Br-Br bond.

---

---


---

Bond energy \_\_\_\_\_ kJ / mole

**(3)**

- (d) **Figure 3** shows the reaction between ethene and chlorine and is similar to the reaction between ethene and bromine.

**Figure 3**



“The more energy levels (shells) of electrons an atom has, the weaker the covalent bonds that it forms.”

Use the above statement to predict and explain how the overall energy change for the reaction of ethene with chlorine will differ from the overall energy change for the reaction of ethene with bromine.

---

---

---

---

---

---

---

---

---

---

---

---


(6)

(Total 14 marks)

**7.**

Methane (CH<sub>4</sub>) is used as a fuel.

- (a) The displayed structure of methane is:



Draw a ring around a part of the displayed structure that represents a covalent bond.

(1)

(b) Why is methane a compound?

Tick (✓) **one** box.

Methane contains atoms of two elements, combined chemically.

Methane is not in the periodic table.

Methane is a mixture of two different elements.


(1)

(c) Methane burns in oxygen.

(i) The diagram below shows the energy level diagram for the complete combustion of methane.

Draw and label arrows on the diagram to show:

- the activation energy
- the enthalpy change,  $\Delta H$ .



(2)

(ii) Complete and balance the symbol equation for the complete combustion of methane.



(2)

(iii) Explain why the **incomplete** combustion of methane is dangerous.

---

---

---

---

(2)

(iv) Explain why, in terms of the energy involved in bond breaking and bond making, the combustion of methane is exothermic.

---

---

---

---

---


---

---

(3)

(d) Methane reacts with chlorine in the presence of sunlight.

The equation for this reaction is:



Some bond dissociation energies are given in the table.

Bond	Bond dissociation energy in kJ per mole
C-H	413
C-Cl	327
Cl-Cl	243
H-Cl	432

(i) Show that the enthalpy change,  $\Delta H$ , for this reaction is  $-103$  kJ per mole.

---

---

---

---


---

---

---

(3)

(ii) Methane also reacts with bromine in the presence of sunlight.



This reaction is less exothermic than the reaction between methane and chlorine.

The enthalpy change,  $\Delta H$ , is  $-45$  kJ per mole.

What is a possible reason for this?

Tick (✓) **one** box.

$\text{CH}_3\text{Br}$  has a lower boiling point than  $\text{CH}_3\text{Cl}$

The C-Br bond is weaker than the C-Cl bond.

The H-Cl bond is weaker than the H-Br bond.

Chlorine is more reactive than bromine.

(1)  
(Total 15 marks)