

Name:

Date:

P3 - Test 1  
PARTICLE MODEL OF MATTER  
Beginner

**GCSE**

**PHYSICS**

**AQA - Triple Science**

Mark

Grade

---

### Materials

For this paper you must have:

- Ruler
- Pencil and Rubber
- Scientific calculator, which you are expected to use when appropriate

### Instructions

- Answer all questions
- Answer questions in the space provided
- All working must be shown


### Information

- The marks for the questions are shown in brackets

1.


(a) The diagrams show the arrangement of the particles in a solid and in a gas.

Each circle represents one particle.



(i) Complete the diagram below to show the arrangement of the particles in a liquid.

**Liquid**



(2)

(ii) Explain, in terms of the particles, why gases are easy to compress.

---

---


---

---

(2)

- (b) The diagram below shows the model that a science teacher used to show her students that there is a link between the temperature of a gas and the speed of the gas particles.

The ball-bearings represent the gas particles. Switching the motor on makes the ball-bearings move around in all directions.



- (i) How is the motion of the ball-bearings similar to the motion of the gas particles?

---

---

(1)

- (ii) The faster the motor runs, the faster the ball-bearings move. Increasing the speed of the motor is like increasing the temperature of a gas.

Use the model to predict what happens to the speed of the gas particles when the temperature of a gas is increased.

---

---

(1)


(Total 6 marks)

2.

A student investigated how the pressure of a gas varied with the volume of the gas.

The mass and temperature of the gas were constant.

The diagram shows the equipment the student used.



(a) What is the range of the syringe?

Tick **one** box.

0 to 1 cm<sup>3</sup>

0 to 5 cm<sup>3</sup>

0 to 20 cm<sup>3</sup>

0 to 25 cm<sup>3</sup>

(1)

(b) What type of variable was the mass of gas?

Tick **one** box.

Control


Dependent

Independent

(1)

The student compressed the gas in the syringe and read the pressure from the pressure gauge.

The graph shows the student's results.



(c) The student concluded that when the pressure was multiplied by the corresponding volume the answer was the same.

Use data from the graph to show that the student's conclusion was correct.

---

---

---

---

---

---

---

(2)

(d) Complete the sentences.

Choose the answers from the box.

Each answer may be used once, more than once or not at all.

<b>decreases</b>	<b>increases</b>	<b>remains the same</b>
------------------	------------------	-------------------------

When the gas is compressed, the volume of gas in the syringe \_\_\_\_\_ .

So the number of collisions each second between the gas particles inside the syringe and the inside surface of the syringe \_\_\_\_\_ .

This means the force exerted on the inside surface of the container walls \_\_\_\_\_ .

(3)

(Total 7 marks)

**3.**

A student wanted to determine the density of a small piece of rock.

(a) Describe how the student could measure the volume of the piece of rock.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(4)

(b) The volume of the piece of rock was 18.0 cm<sup>3</sup>.

The student measured the mass of the piece of rock as 48.6 g.

Calculate the density of the rock in g/cm<sup>3</sup>.

Use the equation:

$$\text{density} = \frac{\text{mass}}{\text{volume}}$$

---

---


---

---

Density = \_\_\_\_\_ g/cm<sup>3</sup>

(2)

The graph below shows the densities of different types of rock.



(c) What is the most likely type of rock that the student had?

Tick **one** box.

Basalt

Flint

Granite

Limestone

Sandstone

(1)

(d) Give **one** source of error that may have occurred when the student measured the volume of the rock.

---

---

(1)

(e) How would the error you described in part **(d)** affect the measured volume of the rock?

---

---

(1)

(Total 9 marks)




4.

A student investigated the cooling effect of evaporation.

She used the equipment (datalogger and probe) shown in **Figure 1** to measure how the temperature of a liquid changed as the liquid evaporated.

**Figure 1**



(a) Which type of variable was the temperature in this investigation?

Tick (✓) **one** box.

	Tick (✓)
control	
dependent	
independent	

(1)

(b) Before the investigation started, the student checked the accuracy of three different temperature probes. The student put the probes in a beaker of boiling water that had a temperature of 100.0 °C.

The readings from the three temperature probes are shown in **Figure 2**.

**Figure 2**

Probe A	Probe B	Probe C
99.8	100.1	103.2

Which **one** of the temperature probes, **A**, **B** or **C**, was **least** accurate?

Write the correct answer in the box.

Give a reason for your answer.


---

---

(2)

(c) **Figure 3** shows how the temperature recorded changed during the investigation.

**Figure 3**



(i) Use **Figure 3** to determine the lowest temperature recorded as the liquid evaporated.

Temperature = \_\_\_\_\_ °C

(1)

(ii) Use **Figure 3** to determine how long it took for all the liquid to evaporate. Give a reason for your answer.

Time = \_\_\_\_\_ seconds

Reason: \_\_\_\_\_

---

(2)

- (iii) How would increasing the starting temperature of the liquid above 20 °C affect the rate of evaporation of the liquid?

---

---

(1)

(Total 7 marks)

5.

The diagram below shows a cyclist riding along a flat road.



- (a) Complete the sentence.

Choose answers from the box.

chemical	elastic potential	gravitational potential	kinetic
----------	-------------------	-------------------------	---------

As the cyclist accelerates, the \_\_\_\_\_ energy store in the cyclist's body decreases and the \_\_\_\_\_ energy of the cyclist increases.

(2)

- (b) The mass of the cyclist is 80 kg. The speed of the cyclist is 12 m/s.

Calculate the kinetic energy of the cyclist.

Use the equation:

$$\text{kinetic energy} = 0.5 \times \text{mass} \times (\text{speed})^2$$

---

---

---

---

Kinetic energy = \_\_\_\_\_ J

(2)

(c) When the cyclist uses the brakes, the bicycle slows down.

This causes the temperature of the brake pads to increase by 50 °C.

The mass of the brake pads is 0.040 kg.

The specific heat capacity of the material of the brake pads is 480 J/kg °C.

Calculate the change in thermal energy of the brake pads.

Use the equation:

change in thermal energy = mass × specific heat capacity × temperature change

---

---

---

Change in thermal energy = \_\_\_\_\_ J

(2)

(d) How is the internal energy of the particles in the brake pads affected by the increase in temperature?

Tick **one** box.

Decreased

Increased

Not affected

(1)

(Total 7 marks)